REFERENCES

LeResche L, Saunders K, Von Korff MR, Barlow W, Dworkin SF (1997). Use of exogenous hormones and risk of temporomandibular disorder pain, Pain, 69, 153-60.

Bonjardim LR, Gavião MB, Pereira LJ, Castelo PM, Garcia RC (2005). Signs and symptoms of temporomandibular disorders in adolescents, Brazilian Oral Research, 19, 93-98.

Garcia AL, Lacerda NJ, Pereira SLS (2001). Evaluation of the degree of dysfunction of the temporomandibular joint and of mandibular movements in young adults, Rev Assoc Paul Cir Dent,51,46-51.

Pedroni CR, De Oliveira AS, Guaratini MI (2003). Prevalence study of signs and symptoms of temporomandibular disorders in university students, Journal of Oral Rehabilitation, 30, 283-289.

Rugh JD & Solberg WK (1985). Oral Health status in United Stated:Temporomandibular Disorders, Journal Of dental Education, 49, 398-405.

Schiffman E, Fricton JR (1998). Epidemiology of TMJ and craniofacial pain: diagnosis and management, In: The TMJ disorders management of the craniomandibular complex. Kraus SL. ed. New York, Churchill Livingstone, 23-38.

Gray RJ, Davies SJ, Quayle AA (1994). A clinical approach to temporomandibular disorders, Brazilian Dental Journal, 176, 429-435.

Bagis B, E A Ayaz, S Turgut, R Durkan, M Ozcan (2012). Gender difference in prevalence of signs and symptoms of temporomandibular joint disorders: a retrospective study on 243 consecutive patients. International journal of medical sciences, 9(7), 539–44

Sessle & Barry J (2014).Newly Developed Diagnostic Criteria for Temporomandibular Disorders, and the Journal of Oral & Facial Pain and Headache, 28, 1, 6-27.

Bonjardim LR, Lopes-Filho RJ**,**Amado G**,** Albuquerque RL**,** Goncalves SR, (2009). Association between symptoms of temporomandibular disorders and gender, morphological occlusion, and psychological factors in a group of university students, Indian Journal of Dental Research, 20(2), 190–4.

Cooper, B.C. & Kleinberg, I(2007). Examination of a large patient population for the presence of symptoms and signs of temporomandibular disorders, Journal of craniomandibular practice, 25(2), 114–26.

Dulcic, N, Panduric J, Simunkovic SK, Badel T, Celic R (2003). Incidence of temporomandibular disorders at tooth loss in the supporting zones, Collegium antropologicum, 27(2), 61–67.

Garcia, AR, Gallo AK, Zuim PR, Dos Santos DM, Antenucci RM(2008). Evaluation of temporomandibular joint noise in partially edentulous patients. Acta odontologica latinoamericana, 21, 21–27.

Gupta, SK, *PK Pratibha PK, KM Bhat KM, S Mutalik S, V Guddattu*(2014). Non-replaced mandibular first molars and temporomandibular joint dysfunction, Nepal Journal of Medical Sciences, 3(1), 57-62.

LeResche L(1997). Epidemiology of temporomandibular disorders: implications for the investigation of etiologic factors. Critical reviews in oral biology and medicine, American Association of Oral Biologists, 8(3), 291–305.

Liu, F & Steinkeler, A(2013). Epidemiology, diagnosis, and treatment of temporomandibular disorders. Dental clinics of North America, 57(3), 465–79.

Shet, R, Rao S, Patel R, Suvvati P, Sadar LR, Yadav RD(2013). Prevalence of temporomandibular joint dysfunction and its signs among the partially edentulous patients in a village of North Gujarat,The Journal of Contemporary Dental Practice, 14,1151–1155.

Nomura K, Vitti M, de Oliveira AS, CC Thais, Semprini M, Siessere S (2007). Use of the Fonseca's Questionnaire to Assess the Prevalence and Severity of Temporomandibular Disorders in Brazilian Dental Undergraduates, Brazilian Dental Journal,18(2), 163-167.

Oliveira AS, Dias EM, Contato RG, Berzin F (2006). Prevalence study of signs and symptoms of temporomandibular disorder in Brazilian college students, Brazilian Oral Research, 20(1), 3-7.

Otuyemi O.D, Ototade F. J, Ugboko VI, Ndukwe KC, Olusile OA (2000). Prevalence of signs and symptoms of temporomandibular disorders in young Nigerian adults, Journal of Orthodontics,27(1), 61-65.

Shiau Y-Y, Chang C (1992). An epidemiological study of temporomandibular disorders in university students of Taiwan. Community Dent Oral Epidemiol, 20, 43-7.

Santos Berni KC, Dibai-Filho AV, Bigaton DR (2015). Accuracy of the Fonseca anamnestic index in the identification of myogenous temporomandibular disorder in female community cases, Journal of Bodywork & Movement Therapies, 19, 404- 409.

Tallents, RH, (2002). Prevalence of missing posterior teeth and intraarticular temporomandibular disorders. The Journal of prosthetic dentistry, 87(1), 45–50.

Wang, MQ, Xue F, He JJ, Chen JH, Chen CS, Raustia A(2009). Missing posterior teeth and risk of temporomandibular disorders. Journal of dental research, 88(10), 942–5.

Martínez CC (2013). Prevalence of temporomandibular disorders according to RDC/TMD, in patients of asub-urban community of Puebla, Mexico, Columbian Journal of Dental Research, 4, 10.

Manfredini D, Chiappe G & Bosco M (2006). Research diagnostic criteria for temporomandibular disorders (RDC/TMD) axis I diagnoses in an Italian patient population. Journal of Oral Rehabilitation, 33(8),551-8.

Yap A, Sworkin S, Chuba E, Tan K & Tan H (2003). Prevalence of temporomandibular disorder subtypes, psychological distress, and psychosocial dysfunction in Asian patients. Journal of Orofacial Pain, 17, 21, 586-587.

Fonseca DM. Disfunção Craniomandibular (1992). Diagnostic pela anamneses, Faculdade de Odontologia de Bauru da USP, 16-23

Bevilaqua D, Chaves T, Oliveira A, Monteiro V (2006). Anamnestic index severity and signs and symptoms of temporomandibular disorder, Craniomandibular, 24, 112–18.

Fonseca DM, Bonfante G, Valle AL, Freitas SF (1994). Diagnóstico pela anamnese da disfunção craniomandibular, 42(1), 23-8.

Berni, Dibai-Filho & Rodrigues-Bigaton (2015). Accuracy of the Fonseca anamnestic index in the identification of myogenous temporomandibular disorder in female community cases, Journal of body work and therapies, 19,404-409.

Wiese M, Svensson P, Bakke M, List T, Hintze H, Petersson A, Knutsson K& Wenzel A (2009).Association between temporomandibular joint symptoms, signs, and clinical diagnosis using the RDC/TMD and radiographic findings in temporomandibular joint tomograms, Oral Surgery Oral Medicine Oral Pathology Oral Radiology Endodontic, 108(3), 117-24.

Fonseca’s questionnaire.

Questions	No	sometimes	Yes
1-Is it hard for you to open your mouth?			
2- Is it hard for you to move your mandible from side to side?			
3- Do you get tired /muscular pain while chewing?			
4 - Do you have frequent headaches?			
5- Do you have pain on the nape or stiff neck?			
6- Do you have earaches or pain in craniomandibular joints?			
7- Have you noticed any TMJ clickingwhile chewing or when you open your mouth?			
8- Do you clench or grind your teeth ?			
9- Do you feel your teeth do not articulate well?			
10- Do you consider yourself a tense (nervous) person?			